

# $\mathcal{PT}$ -symmetric waveguides

Denis Borisov<sup>a,b</sup> and David Krejčířík<sup>b</sup>

a) *Department of Physics and Mathematics, Bashkir State Pedagogical University, October rev. st. 3a, 450000, Ufa, Russia*

b) *Department of Theoretical Physics, Nuclear Physics Institute, Academy of Sciences, 25068 Řež, Czechia*

borisovdi@yandex.ru, krejcirik@ujf.cas.cz

11 November 2008

## Abstract

We introduce a planar waveguide of constant width with non-Hermitian  $\mathcal{PT}$ -symmetric Robin boundary conditions. We study the spectrum of this system in the regime when the boundary coupling function is a compactly supported perturbation of a homogeneous coupling. We prove that the essential spectrum is positive and independent of such perturbation, and that the residual spectrum is empty. Assuming that the perturbation is small in the supremum norm, we show that it gives rise to real weakly-coupled eigenvalues converging to the threshold of the essential spectrum. We derive sufficient conditions for these eigenvalues to exist or to be absent. Moreover, we construct the leading terms of the asymptotic expansions of these eigenvalues and the associated eigenfunctions.

**MSC 2000:** 35P15, 35J05, 47B44, 47B99.

**Keywords:** non-self-adjointness,  $J$ -self-adjointness,  $\mathcal{PT}$ -symmetry, waveguides, Robin boundary conditions, Robin Laplacian, eigenvalue and eigenfunction asymptotics, essential spectrum, reality of the spectrum.

**To appear in:** Integral Equations and Operator Theory

<http://dx.doi.org/10.1007/s00020-008-1634-1>

# 1 Introduction

There are two kinds of motivations for the present work. The first one is due to the growing interest in spectral theory of non-self-adjoint operators. It is traditionally relevant to the study of dissipative processes, resonances if one uses the mathematical tool of complex scaling, and many others. The most recent and conceptually new application is based on the potential quantum-mechanical interpretation of non-Hermitian Hamiltonians which have real spectra and are invariant under a simultaneous  $\mathcal{P}$ -parity and  $\mathcal{T}$ -time reversal. For more information on the subject, we refer to the pioneering work [3] and especially to the recent review [2] with many references.

The other motivation is due to the interesting phenomena of the existence of bound states in quantum-waveguide systems intensively studied for almost two decades. Here we refer to the pioneering work [12] and to the reviews [10, 21]. In these models the Hamiltonian is self-adjoint and the bound states – often without classical interpretations – correspond to an electron trapped inside the waveguide.

In this paper we unify these two fields of mathematical physics by considering a quantum waveguide modelled by a non-Hermitian  $\mathcal{PT}$ -symmetric Hamiltonian. Our main interest is to develop a spectral theory for the Hamiltonian and demonstrate the existence of eigenvalues outside the essential spectrum. For non-self-adjoint operators the location of the various essential spectra is often as much as one can realistically hope for in the absence of the powerful tools available when the operators are self-adjoint, notably the spectral theorem and minimax principle. In the present paper we overcome this difficulty by using perturbation methods to study the point spectrum in the weak-coupling regime. In certain situations we are also able to prove that the total spectrum is real.

Let us now briefly recall the notion of  $\mathcal{PT}$ -symmetry. If the underlying Hilbert space of a Hamiltonian  $H$  is the usual realization of square integrable functions  $L_2(\mathbb{R}^n)$ , the  $\mathcal{PT}$ -symmetry invariance can be stated in terms of the commutator relation

$$(\mathcal{PT})H = H(\mathcal{PT}), \quad (1.1)$$

where the parity and time reversal operators are defined by  $(\mathcal{P}\psi)(x) := \psi(-x)$  and  $\mathcal{T}\psi := \overline{\psi}$ , respectively. In most of the  $\mathcal{PT}$ -symmetric examples  $H$  is the Schrödinger operator  $-\Delta + V$  with a potential  $V$  satisfying (1.1), so that  $H^* = \mathcal{T}H\mathcal{T}$  where  $H^*$  denotes the adjoint of  $H$ . This property is known as the  $\mathcal{T}$ -self-adjointness of  $H$  in the mathematical literature [11], and it is not limited to  $\mathcal{PT}$ -symmetric Schrödinger operators. More generally, given any linear operator  $H$  in an abstract Hilbert space  $\mathcal{H}$ , we understand the  $\mathcal{PT}$ -symmetry property as a special case of the  $J$ -self-adjointness of  $H$ :

$$H^* = JHJ, \quad (1.2)$$

where  $J$  is a conjugation operator, *i.e.*,

$$\forall \phi, \psi \in \mathcal{H}, \quad (J\phi, J\psi)_{\mathcal{H}} = (\psi, \phi)_{\mathcal{H}}, \quad J^2\psi = \psi.$$

This setting seems to be adequate for a rigorous formulation of  $\mathcal{PT}$ -symmetric problems, and alternative to that based on Krein spaces [22, 24].

The nice feature of the property (1.2) is that  $H$  “is not too far” from the class of self-adjoint operators. In particular, the eigenvalues are found to be

real for many  $\mathcal{PT}$ -symmetric Hamiltonians [28, 9, 22, 8, 26, 7, 20]. However, the situation is much less studied in the case when the resolvent of  $H$  is not compact.

The spectral analysis of non-self-adjoint operators is more difficult than in the self-adjoint case, partly because the residual spectrum is in general not empty for the former. One of the goals of the present paper is to point out that the existence of this part of spectrum is always ruled out by (1.2):

**Fact.** *Let  $H$  be a densely defined closed linear operator in a Hilbert space satisfying (1.2). Then the residual spectrum of  $H$  is empty.*

The proof follows easily by noticing that the kernels of  $H - \lambda$  and  $H^* - \bar{\lambda}$  have the same dimension [11, Lem. III.5.4] and by the general fact that the orthogonal complement of the range of a densely defined closed operator in a Hilbert space is equal to the kernel of its adjoint. The above result is probably not well known in the  $\mathcal{PT}$ -symmetry community.

We continue with an informal presentation of our model and main spectral results obtained in this paper. The rigorous and more detailed statements are postponed until the next section because they require a number of technical definitions.

The Hamiltonian we consider in this paper acts as the Laplacian in the Hilbert space of square integrable functions over a straight planar strip and the non-Hermiticity enters through  $\mathcal{PT}$ -symmetric boundary conditions only. The boundary conditions are of Robin type but with imaginary coupling. The  $\mathcal{PT}$ -symmetric invariance then implies that we actually deal with an electromagnetic waveguide with radiation/dissipative boundary conditions. In fact, the one-dimensional spectral problem in the waveguide cross-section has been studied recently in [20] (see also [19]) and our model can be viewed as a two-dimensional extension of the former.

Schrödinger-type operators with similar non-Hermitian boundary conditions were studied previously by Kaiser, Neidhardt and Rehberg [17, 16, 15]. In their papers, motivated by the needs of semiconductor physics, the configuration space is a bounded domain and the boundary coupling function is such that the Hamiltonian is a dissipative operator. The latter excludes the  $\mathcal{PT}$ -symmetric models of [20] and the present paper.

The  $\mathcal{T}$ -self-adjointness property (1.2) of our Hamiltonian is proved in Section 3. If the boundary coupling function is constant, the spectral problem can be solved by separation of variables and we find that the spectrum is purely essential, given by a positive semibounded interval (*cf* Section 4). In Section 5 we prove that the essential spectrum is stable under compactly supported perturbations of the coupling function. Consequently, the essential spectrum is always real in our setting, however, it exhibits important differences as regards similar self-adjoint problems. Namely, it becomes as a set independent of the value of the coupling function at infinity when the latter overpasses certain critical value.

In Section 6 we study the point spectrum. We focus on the existence of eigenvalues emerging from the threshold of the essential spectrum in the limit when the compactly supported perturbation of the coupling function tends to zero in the supremum norm. It turns out that the weakly-coupled eigenvalues may or may not exist, depending on mean values of the local perturbation. In

the case when the point spectrum exists, we derive asymptotic expansions of the eigenvalues and the associated eigenfunctions.

Because of the singular nature of the  $\mathcal{PT}$ -symmetric interaction, our example is probably the simplest non-trivial, multidimensional  $\mathcal{PT}$ -symmetric model whatsoever for which both the point and essential spectra exist. We hope that the present work will stimulate more research effort in the direction of spectral and scattering properties of the present and other non-Hermitian  $\mathcal{PT}$ -symmetric operators.

## 2 Main results

Given a positive number  $d$ , we write  $I := (0, d)$  and consider an infinite straight strip  $\Omega := \mathbb{R} \times I$ . We split the variables consistently by writing  $x = (x_1, x_2)$  with  $x_1 \in \mathbb{R}$  and  $x_2 \in I$ . Let  $\alpha$  be a bounded real-valued function on  $\mathbb{R}$ ; occasionally we shall denote by the same symbol the function  $x \mapsto \alpha(x_1)$  on  $\Omega$ . The object of our interest is the operator in the Hilbert space  $L_2(\Omega)$  which acts as the Laplacian and satisfies the following  $\mathcal{PT}$ -symmetric boundary conditions:

$$\partial_2 \Psi + i\alpha \Psi = 0 \quad \text{on } \partial\Omega. \quad (2.1)$$

More precisely, we introduce

$$H_\alpha \Psi := -\Delta \Psi, \quad \Psi \in \mathfrak{D}(H_\alpha) := \{\Psi \in W_2^2(\Omega) \mid \Psi \text{ satisfies (2.1)}\}, \quad (2.2)$$

where the action of  $H_\alpha$  should be understood in the distributional sense and (2.1) should be understood in the sense of traces [1]. In Section 3 we show that  $H_\alpha$  is well defined in the sense that it is an  $m$ -sectorial operator and that its adjoint is easy to identify:

**Theorem 2.1.** *Let  $\alpha \in W_\infty^1(\mathbb{R})$ . Then  $H_\alpha$  is an  $m$ -sectorial operator in  $L_2(\Omega)$  satisfying*

$$H_\alpha^* = H_{-\alpha}. \quad (2.3)$$

Of course,  $H_\alpha$  is not self-adjoint unless  $\alpha$  vanishes identically (in this case  $H_0$  is the Neumann Laplacian in  $L_2(\Omega)$ ). However,  $H_\alpha$  is  $\mathcal{T}$ -self-adjoint, *i.e.*, it satisfies (1.2) with  $J$  being the complex conjugation  $\mathcal{T} : \Psi \mapsto \bar{\Psi}$ . Indeed,  $H_\alpha$  satisfies the relation (2.3) and it is easy to see that

$$H_{-\alpha} = \mathcal{T} H_\alpha \mathcal{T}. \quad (2.4)$$

This reflects the  $\mathcal{PT}$ -symmetry (1.1) of our problem, with  $\mathcal{P}$  being defined by  $(\mathcal{P}\Psi)(x) := \Psi(x_1, d - x_2)$ .

An important property of an operator  $H$  in a Hilbert space  $\mathcal{H}$  being  $m$ -sectorial is that it is closed. Then, in particular, the spectrum  $\sigma(H)$  is well defined as the set of complex points  $z$  such that  $H - z$  is not bijective as the operator from  $\mathfrak{D}(H)$  to  $\mathcal{H}$ . Furthermore, its spectrum is contained in a sector of complex numbers  $z$  such that  $|\arg(z - \gamma)| \leq \theta$  with some  $\gamma \in \mathbb{R}$  and  $\theta \in [0, \pi/2)$ . In our case, however, we are able to establish a stronger result

$$\sigma(H_\alpha) \subseteq \Xi_\alpha := \left\{ z \in \mathbb{C} : \operatorname{Re} z \geq 0, |\operatorname{Im} z| \leq 2 \|\alpha\|_{L_\infty(\mathbb{R})} \sqrt{\operatorname{Re} z} \right\}. \quad (2.5)$$

This follows directly from Lemma 3.1 on which the proof of Theorem 2.1 is based (*cf* the end of Section 3 for more details). Consequently, the resolvent set  $\rho(H_\alpha) := \mathbb{C} \setminus \sigma(H_\alpha)$  contains the complement of  $\Xi_\alpha$  and we have the bound

$$\|(H_\alpha - z)^{-1}\| \leq 1/\text{dist}(z, \partial\Xi_\alpha) \quad \text{for all } z \in \mathbb{C} \setminus \Xi_\alpha, \quad (2.6)$$

where  $\|\cdot\|$  denotes the operator norm in  $L_2(\Omega)$ .

Given a closed operator  $H$  in a Hilbert space  $\mathcal{H}$ , we use the following decomposition of the spectrum  $\sigma(H)$ :

**Definition.** The *point* spectrum  $\sigma_p(H)$  equals the set of points  $\lambda$  such that  $H - \lambda$  is not injective. The *essential* spectrum  $\sigma_e(H)$  equals the set of points  $\lambda$  such that  $H - \lambda$  is not Fredholm. Finally, the *residual* spectrum  $\sigma_r(H)$  equals the set of points  $\lambda$  such that  $H - \lambda$  is injective but the range of  $H - \lambda$  is not dense in  $\mathcal{H}$ .

*Remark 2.1.* 1. The reader is warned that various other types of essential spectra of non-self-adjoint operators are used in the literature; *cf* [11, Chapt. IX] for five distinct definitions and a detailed description of their properties. Among them we choose that of Wolf [27], which is in general larger than that of Kato [18, Sec. IV.5.6] based on violating the semi-Fredholm property. (Recall that a closed operator in a Hilbert space is called Fredholm if its range is closed and both its kernel and its cokernel are finite-dimensional, while it is called semi-Fredholm if its range is closed and its kernel or its cokernel is finite-dimensional.) However, since our operator  $H_\alpha$  is  $\mathcal{T}$ -self-adjoint, the majority of the different definitions coincide [11, Thm IX.1.6], in particular the two above, and that is why we use the common notation  $\sigma_e(\cdot)$  in this paper. Then our choice also coincides with the definition of “continuous spectrum” as used for instance in the Glazman’s book [14].

2. We indeed have the decomposition (*cf* [14, Sec. I.1.1])

$$\sigma(H) = \sigma_p(H) \cup \sigma_e(H) \cup \sigma_r(H),$$

but note that there might be intersections on the right hand side. In particular,  $\sigma_e(H)$  contains eigenvalues of infinite geometric multiplicity.

3. On the other hand, the definitions of point and residual spectra are standard and they form disjoint subsets of  $\sigma(H)$ . Recalling the general fact [18, Sec. V.3.1] that the orthogonal complement of the range of a densely defined closed operator in a Hilbert space is equal to the kernel of its adjoint, we obtain the following characterization of the residual spectrum in terms of the point spectrum of the operator and its adjoint:

$$\sigma_r(H) = \{\lambda \in \mathbb{C} \mid \bar{\lambda} \in \sigma_p(H^*) \text{ \& } \lambda \notin \sigma_p(H)\}. \quad (2.7)$$

The  $\mathcal{T}$ -self-adjointness of  $H_\alpha$  immediately implies:

**Corollary 2.1.** *Suppose the hypothesis of Theorem 2.1. Then*

$$\sigma_r(H_\alpha) = \emptyset.$$

*Proof.* We repeat the proof sketched in Introduction. Since  $H_\alpha$  is  $\mathcal{T}$ -self-adjoint, it is easy to see that  $\lambda$  is an eigenvalue of  $H_\alpha$  (with eigenfunction  $\Psi$ ) if, and only if,  $\bar{\lambda}$  is an eigenvalue of  $H_\alpha^*$  (with eigenfunction  $\bar{\Psi}$ ). It is then clear from the general identity (2.7) that the residual spectrum of  $H_\alpha$  must be empty.  $\square$

The case of uniform boundary conditions, *i.e.* when  $\alpha$  equals identically a constant  $\alpha_0$ , can be solved by separation of variables (*cf* Section 4). We find

$$\sigma(H_{\alpha_0}) = \sigma_e(H_{\alpha_0}) = [\mu_0^2, +\infty), \quad (2.8)$$

where the threshold  $\mu_0^2$ , with the notation

$$\mu_0 = \begin{cases} \alpha_0 & \text{if } |\alpha_0| \leq \pi/d, \\ \pi/d & \text{if } |\alpha_0| > \pi/d, \end{cases} \quad (2.9)$$

denotes the bottom of the spectrum of the “transverse” operator

$$\begin{aligned} -\Delta_{\alpha_0}^I \psi &:= -\psi'', \\ \psi \in \mathfrak{D}(-\Delta_{\alpha_0}^I) &:= \{\psi \in W_2^2(I) \mid \psi' + i\alpha_0 \psi = 0 \text{ at } \partial I\}. \end{aligned} \quad (2.10)$$

The operator  $-\Delta_{\alpha_0}^I$  was studied in [20]. Its spectrum is purely discrete and real:

$$\sigma(-\Delta_{\alpha_0}^I) = \{\mu_j^2\}_{j=0}^\infty, \quad (2.11)$$

where  $\mu_0$  has been introduced in (2.9),

$$\mu_1 = \begin{cases} \alpha_0 & \text{if } |\alpha_0| > \pi/d, \\ \pi/d & \text{if } |\alpha_0| \leq \pi/d, \end{cases} \quad \text{and} \quad \mu_j := \pi j/d \quad \text{for } j \geq 2.$$

Making the hypothesis

$$\alpha_0 d/\pi \notin \mathbb{Z} \setminus \{0\}, \quad (2.12)$$

the eigenvalues of  $-\Delta_{\alpha_0}^I$  are simple and the corresponding set of eigenfunctions  $\{\psi_j\}_{j=0}^\infty$  can be chosen as

$$\psi_j(x_2) := \cos(\mu_j x_2) - i \frac{\alpha_0}{\mu_j} \sin(\mu_j x_2). \quad (2.13)$$

We refer to Section 4.1 for more results about the operator  $-\Delta_{\alpha_0}^I$ .

Let us now turn to the non-trivial case of variable coupling function  $\alpha$ . Among a variety of possible situations, in this paper we restrict the considerations to local perturbations of the uniform case. Namely, we always assume that the difference  $\alpha - \alpha_0$  is compactly supported.

First of all, in Section 5 we show that the essential component of the spectrum of  $H_\alpha$  is stable under the local perturbation of the uniform case:

**Theorem 2.2.** *Let  $\alpha - \alpha_0 \in C_0(\mathbb{R}) \cap W_\infty^1(\mathbb{R})$  with  $\alpha_0 \in \mathbb{R}$ . Then*

$$\sigma_e(H_\alpha) = [\mu_0^2, +\infty).$$

Notice that the essential spectrum as a set is independent of  $\alpha_0$  as long as  $|\alpha_0| \geq \pi/d$ . This is a consequence of the fact that our Hamiltonian is not Hermitian. On the other hand, it follows that the essential spectrum is real. Recall that the residual spectrum is always empty due to Corollary 2.1. We do not have the proof of the reality for the point spectrum, except for the particular case treated in the next statement:

**Theorem 2.3.** *Let  $\alpha \in C_0(\mathbb{R}) \cap W_\infty^1(\mathbb{R})$  be an odd function. Then*

$$\sigma_p(H_\alpha) \subset \mathbb{R}.$$

Summing up, under the hypotheses of this theorem the total spectrum is real (and in fact non-negative due to (2.5)).

The next part of our results concerns the behavior of the point spectrum of  $H_\alpha$  under a small perturbation of  $\alpha_0$ . Namely, we consider the local perturbation of the form

$$\alpha(x_1) = \alpha_0 + \varepsilon \beta(x_1), \quad (2.14)$$

where  $\beta \in C_0^2(\mathbb{R})$  and  $\varepsilon$  is a small positive parameter. In accordance with Theorem 2.2, in this case the essential spectrum of  $H_\alpha$  coincides with  $[\mu_0^2, +\infty)$ , and this is also the spectrum of  $H_{\alpha_0}$ . Our main interest is focused on the existence and asymptotic behavior of the eigenvalues emerging from the threshold  $\mu_0^2$  due to the perturbation of  $H_{\alpha_0}$  by  $\varepsilon\beta$ .

First we show that the asymptotically Neumann case is in some sense exceptional:

**Theorem 2.4.** *Suppose  $\alpha_0 = 0$ . Let  $\alpha$  be given by (2.14), where  $\beta \in C_0^2(\mathbb{R})$ . Then the operator  $H_\alpha$  has no eigenvalues converging to  $\mu_0^2$  as  $\varepsilon \rightarrow +0$ .*

The problem of existence of the weakly-coupled eigenvalues is more subtle as long as  $\alpha_0 \neq 0$ . To present our results in this case, we introduce an auxiliary sequence of functions  $v_j : \mathbb{R} \rightarrow \mathbb{R}$  by

$$v_j(x_1) := \begin{cases} -\frac{1}{2} \int_{\mathbb{R}} |x_1 - t_1| \beta(t_1) dt_1 & \text{if } j = 0, \\ \frac{1}{2\sqrt{\mu_j^2 - \mu_0^2}} \int_{\mathbb{R}} e^{-\sqrt{\mu_j^2 - \mu_0^2}|x_1 - t_1|} \beta(t_1) dt_1 & \text{if } j \geq 1. \end{cases} \quad (2.15)$$

Denoting  $\langle f \rangle = \int_{\mathbb{R}} f(x_1) dx_1$  for any  $f \in L_1(\mathbb{R})$ , we introduce a constant  $\tau$ , depending on  $\beta$ ,  $d$  and  $\alpha_0$ , by

$$\tau := \begin{cases} 2\alpha_0^2 \langle \beta v_0 \rangle + \frac{2\alpha_0}{d} \sum_{j=1}^{\infty} \frac{\mu_j^2 \langle \beta v_j \rangle}{\mu_j^2 - \mu_0^2} \tan \frac{\alpha_0 d + j\pi}{2} & \text{if } |\alpha_0| < \frac{\pi}{d}, \\ \frac{2\alpha_0 \pi^2 \cot \frac{\alpha_0 d}{2}}{(\mu_1^2 - \mu_0^2) d^3} \langle \beta v_1 \rangle + \frac{8\pi^2}{(\mu_1^2 - \mu_0^2) d^4} \sum_{j=1}^{\infty} \frac{\mu_{2j}^2 \langle \beta v_{2j} \rangle}{\mu_{2j}^2 - \mu_1^2} & \text{if } |\alpha_0| > \frac{\pi}{d}. \end{cases}$$

It will be shown in Section 6.3 that the series converge. Finally, we denote  $\Omega_a := \Omega \cap \{x : |x_1| < a\}$  for any positive  $a$ . Now we are in a position to state our main results about the point spectrum.

**Theorem 2.5.** *Suppose  $|\alpha_0| < \pi/d$ . Let  $\alpha$  be given by (2.14), where  $\beta \in C_0^2(\mathbb{R})$ .*

1. *If  $\alpha_0 \langle \beta \rangle < 0$ , there exists the unique eigenvalue  $\lambda_\varepsilon$  of  $H_\alpha$  converging to  $\mu_0^2$  as  $\varepsilon \rightarrow +0$ . This eigenvalue is simple and real, and satisfies the asymptotic formula*

$$\lambda_\varepsilon = \mu_0^2 - \varepsilon^2 \alpha_0^2 \langle \beta \rangle^2 + 2\varepsilon^3 \alpha_0 \tau \langle \beta \rangle + \mathcal{O}(\varepsilon^4).$$

*The associated eigenfunction  $\Psi_\varepsilon$  can be chosen so that it satisfies the asymptotics*

$$\Psi_\varepsilon(x) = \psi_0(x_2) + \mathcal{O}(\varepsilon) \quad (2.16)$$

in  $W_2^2(\Omega_a)$  for each  $a > 0$ , and behaves at infinity as

$$\Psi_\varepsilon(x) = e^{-\sqrt{\mu_0^2 - \lambda_\varepsilon}|x_1|} \psi_0(x_2) + \mathcal{O}(e^{-\sqrt{\mu_0^2 - \lambda_\varepsilon}|x_1|}), \quad |x_1| \rightarrow +\infty. \quad (2.17)$$

2. If  $\alpha_0 \langle \beta \rangle > 0$ , the operator  $H_\alpha$  has no eigenvalues converging to  $\mu_0^2$  as  $\varepsilon \rightarrow +0$ .
3. If  $\langle \beta \rangle = 0$ , and  $\tau > 0$ , there exists the unique eigenvalue  $\lambda_\varepsilon$  of  $H_\alpha$  converging to  $\mu_0^2$  as  $\varepsilon \rightarrow +0$ . This eigenvalue is simple and real, and satisfies the asymptotics

$$\lambda_\varepsilon = \mu_0^2 - \varepsilon^4 \tau^2 + \mathcal{O}(\varepsilon^5). \quad (2.18)$$

The associated eigenfunction can be chosen so that the relations (2.16) and (2.17) hold true.

4. If  $\langle \beta \rangle = 0$ , and  $\tau < 0$ , the operator  $H_\alpha$  has no eigenvalues converging to  $\mu_0^2$  as  $\varepsilon \rightarrow +0$ .

**Theorem 2.6.** Suppose  $|\alpha_0| > \pi/d$  and (2.12). Let  $\alpha$  be given by (2.14) where  $\beta \in C_0^2(\mathbb{R})$ .

1. If  $\tau > 0$ , there exists the unique eigenvalue  $\lambda_\varepsilon$  of  $H_\alpha$  converging to  $\mu_0$  as  $\varepsilon \rightarrow +0$ , it is simple and real, and satisfies the asymptotics (2.18). The associated eigenfunction can be chosen so that it obeys (2.16) and (2.17).
2. If  $\tau < 0$ , the operator  $H_\alpha$  has no eigenvalues converging to  $\mu_0^2$  as  $\varepsilon \rightarrow +0$ .

In accordance with Theorem 2.5, in the case  $|\alpha_0| < \pi/d$  the existence of a weakly-coupled eigenvalue is determined by the sign of the constant  $\alpha_0$  and that of the mean value of  $\beta$ . In the language of Schrödinger operators (treating  $\alpha$  as a singular potential), it means that a given non-trivial  $\beta$  plays the role of an effective interaction, attractive or repulsive depending upon the sign of  $\alpha_0$ . It is instructive to compare this situation with a self-adjoint waveguide [6], where a similar effective interaction is induced by a local deformation of the boundary. If the boundary is deformed “outward in the mean”, a weakly-coupled bound state exists, while it is absent if the deformation is “inward-pointing in the mean”.

As usual, the critical situation  $\langle \beta \rangle = 0$  is much harder to treat. In our case, one has to check the sign of  $\tau$  to decide whether a weakly-coupled bound state exists. However, it can be difficult to sum up the series in the definition of  $\tau$ . This is why in our next statement we provide a sufficient condition guaranteeing that  $\tau > 0$ .

**Proposition 2.1.** Suppose  $0 < |\alpha_0| < \pi/d$ . Let  $\alpha$  be given by (2.14) where  $\beta(x_1) = \tilde{\beta}(x_1/l)$ ,  $\tilde{\beta} \in C_0^2(\mathbb{R})$ ,  $\langle \tilde{\beta} \rangle = 0$ ,  $l > 0$ . If

$$\left\| \int_{\mathbb{R}} \text{sign}(\cdot - t_1) \tilde{\beta}(t_1) dt_1 \right\|_{L_2(\mathbb{R})}^2 \geq \frac{4 \cot \frac{\alpha_0 d}{2}}{l^2 \alpha_0 d} \left[ \frac{\mu_1^2}{(\mu_1^2 - \mu_0^2)^2} + \frac{d^2}{16\pi^2} + \frac{d^2}{48} \right] \|\tilde{\beta}\|_{L_2(\mathbb{R})}^2,$$

then  $\tau > 0$ .

The meaning of this proposition is that for each positive  $|\alpha_0| < \pi/d$  the perturbation  $\varepsilon\beta$  in the critical regime  $\langle \beta \rangle = 0$  produces a weakly-coupled eigenvalue near the threshold of the essential spectrum provided that the support of  $\beta$

is wide enough. This is in perfect agreement with the critical situation of [6]; according to higher-order asymptotics derived in [5], here the weakly-coupled bound state exists if, and only if, the critical boundary deformation is smeared enough.

In the case  $|\alpha_0| > \pi/d$  a sufficient condition guaranteeing  $\tau > 0$  is given in

**Proposition 2.2.** *Suppose  $|\alpha_0| > \pi/d$  and (2.12). Let  $\alpha$  be given by (2.14) where  $\beta \in C_0^2(\mathbb{R})$ . Let  $m$  be the maximal positive integer such that  $\mu_{2m} < |\alpha_0|$ . If*

$$\alpha_0 \langle \beta v_1 \rangle \cot \frac{\alpha_0 d}{2} \geq \frac{4}{d} \sum_{j=1}^m \frac{\mu_{2j}^2 \langle \beta v_{2j} \rangle}{\mu_1^2 - \mu_{2j}^2}, \quad (2.19)$$

then  $\tau > 0$ .

In Section 6.6 we will show that the inequality (2.19) makes sense. Namely, it will be proved that there exists  $\beta$  such that this inequality holds true, provided that  $\alpha_0$  is close enough to  $\mu_2$  but greater than this value.

*Remark 2.2.* It is useful to make the hypothesis (2.12), since it implies that the “transverse” eigenfunctions (2.13) form a basis (cf (4.1)) and makes it therefore possible to obtain a relatively simple decomposition of the resolvent of  $H_{\alpha_0}$  (cf Lemma 4.3). However, it is rather a technical hypothesis for many of the spectral results (e.g., Theorem 2.2). On the other hand, it seems that the hypothesis is rather crucial for the statement of Theorem 2.6 and Proposition 2.2.

If the hypothesis (2.12) is omitted and  $\alpha_0 = \pi\ell/d$ , with  $\ell \in \mathbb{Z} \setminus \{0\}$ , the threshold of the essential spectrum is  $\pi^2/d^2$ . This point corresponds to a simple eigenvalue of the “transverse” operator  $-\Delta_{\alpha_0}^I$  only if  $|\ell| > 1$ , while it is a double eigenvalue if  $|\ell| = 1$ . Under the hypothesis of Theorems 2.5 and 2.6, the threshold is always a simple eigenvalue of  $-\Delta_{\alpha_0}^I$ , and the proof of the theorems actually employs some sort of “non-degenerate” perturbation theory. In view of this, we conjecture that in the degenerate case  $|\ell| = 1$  two simple eigenvalues (possibly forming a complex conjugate pair) or one double (real) eigenvalue can emerge from the threshold of the essential spectrum for a suitable choice of  $\beta$ , while in the case  $|\ell| > 1$  there can be at most one simple emerging eigenvalue. The question on the asymptotic behaviour of these eigenvalues constitutes an interesting open problem.

### 3 Definition of the operator

In this section we prove Theorem 2.1. Our method is based on the theory of sectorial sesquilinear forms [18, Sec. VI].

In the beginning we assume only that  $\alpha$  is bounded. Let  $h_\alpha$  be the sesquilinear form defined in  $L_2(\Omega)$  by the domain  $\mathfrak{D}(h_\alpha) := W_2^1(\Omega)$  and the prescription  $h_\alpha := h_\alpha^1 + ih_\alpha^2$  with

$$\begin{aligned} h_\alpha^1(\Psi, \Phi) &:= \int_{\Omega} \nabla \Psi(x) \cdot \overline{\nabla \Phi(x)} \, dx, \\ h_\alpha^2(\Psi, \Phi) &:= \int_{\mathbb{R}} \alpha(x_1) \Psi(x_1, d) \overline{\Phi(x_1, d)} \, dx_1 - \int_{\mathbb{R}} \alpha(x_1) \Psi(x_1, 0) \overline{\Phi(x_1, 0)} \, dx_1, \end{aligned}$$

for any  $\Psi, \Phi \in \mathfrak{D}(h_\alpha)$ . Here the dot denotes the scalar product in  $\mathbb{R}^2$  and the boundary terms should be understood in the sense of traces [1]. We write

$h_\alpha[\Psi] := h_\alpha(\Psi, \Psi)$  for the associated quadratic form, and similarly for  $h_\alpha^1$  and  $h_\alpha^2$ .

Clearly,  $h_\alpha$  is densely defined. It is also clear that the real part  $h_\alpha^1$  is a densely defined, symmetric, positive, closed sesquilinear form (it is associated to the self-adjoint Neumann Laplacian in  $L_2(\Omega)$ ). Of course,  $h_\alpha$  itself is not symmetric unless  $\alpha$  vanishes identically; however, it can be shown that it is sectorial and closed. To see it, one can use the perturbation result [18, Thm. VI.1.33] stating that the sum of a sectorial closed form with a relatively bounded form is sectorial and closed provided the relative bound is less than one. In our case, the imaginary part  $h_\alpha^2$  plays the role of the small perturbation of  $h_\alpha^1$  by virtue of the following result.

**Lemma 3.1.** *Let  $\alpha \in L^\infty(\mathbb{R})$ . Then  $h_\alpha^2$  is relatively bounded with respect to  $h_\alpha^1$ , with*

$$|h_\alpha^2[\Psi]| \leq 2 \|\alpha\|_{L^\infty(\mathbb{R})} \|\Psi\|_{L_2(\Omega)} \sqrt{h_\alpha^1[\Psi]} \leq \delta h_\alpha^1[\Psi] + \delta^{-1} \|\alpha\|_{L^\infty(\mathbb{R})}^2 \|\Psi\|_{L_2(\Omega)}^2$$

for all  $\Psi \in W_2^1(\Omega)$  and any positive number  $\delta$ .

*Proof.* By density [1, Thm. 3.18], it is sufficient to prove the inequality for restrictions to  $\Omega$  of functions  $\Psi$  in  $C_0^\infty(\mathbb{R}^2)$ . Then we have

$$|h_\alpha^2[\Psi]| = \left| \int_\Omega \alpha(x_1) \frac{\partial |\Psi(x)|^2}{\partial x_2} dx \right| \leq 2 \|\alpha\|_{L^\infty(\mathbb{R})} \|\Psi\|_{L_2(\Omega)} \|\partial_2 \Psi\|_{L_2(\Omega)},$$

which gives the first inequality after applying  $\|\partial_2 \Psi\|_{L_2(\Omega)} \leq \|\nabla \Psi\|_{L_2(\Omega)}$ . The second inequality then follows at once by means of the Cauchy inequality with  $\delta$ .  $\square$

In view of the above properties, Theorem VI.1.33 in [18], and the first representation theorem [18, Thm. VI.2.1], there exists the unique  $m$ -sectorial operator  $\tilde{H}_\alpha$  in  $L_2(\Omega)$  such that  $h_\alpha(\Psi, \Phi) = (\tilde{H}_\alpha \Psi, \Phi)$  for all  $\Psi \in \mathfrak{D}(\tilde{H}_\alpha) \subset \mathfrak{D}(h_\alpha)$  and  $\Phi \in \mathfrak{D}(h_\alpha)$ , where

$$\mathfrak{D}(\tilde{H}_\alpha) = \{ \Psi \in W_2^1(\Omega) \mid \exists F \in L_2(\Omega), \forall \Phi \in W_2^1(\Omega), h_\alpha(\Psi, \Phi) = (F, \Phi)_{L_2(\Omega)} \}.$$

By integration by parts, it is easy to check that if  $\Psi \in \mathfrak{D}(H_\alpha)$ , it follows that  $\Psi \in \mathfrak{D}(\tilde{H}_\alpha)$  with  $F = -\Delta \Psi$ . That is,  $\tilde{H}_\alpha$  is an extension of  $H_\alpha$  as defined in (2.2). It remains to show that actually  $H_\alpha = \tilde{H}_\alpha$  in order to prove Theorem 2.1. However, the other inclusion holds as a direct consequence of the representation theorem and the following result.

**Lemma 3.2.** *Let  $\alpha \in W_\infty^1(\mathbb{R})$ . For each  $F \in L_2(\Omega)$ , a generalized solution  $\Psi$  to the problem*

$$\begin{cases} -\Delta \Psi = F & \text{in } \Omega, \\ \partial_2 \Psi + i\alpha \Psi = 0 & \text{on } \partial\Omega, \end{cases} \quad (3.1)$$

*belongs to  $\mathfrak{D}(H_\alpha)$ .*

*Proof.* For any function  $\Psi \in W_2^1(\Omega)$ , we introduce the difference quotient

$$\Psi_\delta(x) := \frac{\Psi(x_1 + \delta, x_2) - \Psi(x)}{\delta},$$

where  $\delta$  is a small real number. By standard arguments [23, Ch. III, Sec. 3.4, Thm. 3], the estimate

$$\|\Psi_\delta\|_{L_2(\Omega)} \leq \|\Psi\|_{W_2^1(\Omega)} \quad (3.2)$$

holds true for all  $\delta$  small enough. If  $\Psi$  is a generalized solution to (3.1), then  $\Psi_\delta$  is a generalized solution to the problem

$$\begin{cases} -\Delta \Psi_\delta = F_\delta & \text{in } \Omega, \\ \partial_2 \Psi_\delta + i\alpha \Psi_\delta = g & \text{on } \partial\Omega, \end{cases}$$

where  $g$  denotes the trace of the function  $x \mapsto -i\alpha_\delta(x_1)\Psi(x_1 + \delta, x_2)$  to the boundary  $\partial\Omega$ . Using the “integration-by-parts” formula for the difference quotients,  $(F_\delta, \Phi)_{L_2(\Omega)} = -(F, \Phi_{-\delta})_{L_2(\Omega)}$ , the integral identity corresponding to the weak formulation of the boundary value problem for  $\Psi_\delta$  can be written as follows

$$\begin{aligned} h_\alpha(\Psi_\delta, \Phi) = & -(F, \Phi_{-\delta})_{L_2(\Omega)} - i \int_{\mathbb{R}} \alpha_\delta(x_1) \Psi(x_1 + \delta, d) \overline{\Phi(x_1, d)} dx_1 \\ & + i \int_{\mathbb{R}} \alpha_\delta(x_1) \Psi(x_1 + \delta, 0) \overline{\Phi(x_1, 0)} dx_1, \end{aligned}$$

where  $\Phi \in W_2^1(\Omega)$  is arbitrary. Letting  $\Phi = \Psi_\delta$ , and using the embedding of  $W_2^1(\Omega)$  in  $L_2(\partial\Omega)$ , the boundedness of  $\alpha_\delta$ , Lemma 3.1 and (3.2), the above identity yields

$$\|\Psi_\delta\|_{W_2^1(\Omega)} \leq C,$$

where the constant  $C$  is independent of  $\delta$ . Employing this estimate and proceeding as in the proof of Item b) of Theorem 3 in [23, Ch. III, Sec. 3.4], one can show easily that  $\partial_1 \Psi \in W_2^1(\Omega)$ . Hence,  $\partial_{11} \Psi \in L_2(\Omega)$  and  $\partial_{12} \Psi \in L_2(\Omega)$ .

It follows from standard elliptic regularity theorems (see, *e.g.*, [23, Ch. IV, Sec. 2.2]) that  $\Psi \in W_{2,loc}^2(\Omega)$ . Hence, the first of the equations in (3.1) holds true a.e. in  $\Omega$ . Thus,  $\partial_{22} \Psi = -F - \partial_{11} \Psi \in L_2(\Omega)$ , and therefore  $\Psi \in W_2^2(\Omega)$ .

It remains to check the boundary condition for  $\Psi$ . Integrating by parts, one has

$$\begin{aligned} (F, \Phi)_{L_2(\Omega)} = & h_\alpha(\Psi, \Phi) = (-\Delta \Psi, \Phi)_{L_2(\Omega)} \\ & + \int_{\mathbb{R}} [\partial_2 \Psi(x_1, d) + i\alpha(x_1) \Psi(x_1, d)] \overline{\Phi(x_1, d)} dx_1 \\ & - \int_{\mathbb{R}} [\partial_2 \Psi(x_1, 0) + i\alpha(x_1) \Psi(x_1, 0)] \overline{\Phi(x_1, 0)} dx_1 \end{aligned}$$

for any  $\Phi \in W_2^1(\Omega)$ . This implies the boundary conditions because  $-\Delta \Psi = F$  a.e. in  $\Omega$  and  $\Phi$  is arbitrary.  $\square$

Summing up the results of this section, we get

**Proposition 3.1.** *Let  $\alpha \in W_\infty^1(\mathbb{R})$ . Then  $\tilde{H}_\alpha = H_\alpha$ .*

Theorem 2.1 follows as a corollary of this proposition. In particular, the latter implies that  $H_\alpha$  is  $m$ -sectorial. Moreover, by the first representation theorem, we know that the adjoint  $\tilde{H}_\alpha^*$  is simply obtained as the operator associated with  $h_\alpha^* = h_{-\alpha}$ . This together with Proposition 3.1 proves (2.3).

Let us finally comment on the results (2.5) and (2.6). As a direct consequence of the first inequality of Lemma 3.2, we get that the numerical range of  $H_\alpha (= \tilde{H}_\alpha)$ , defined as the set of all complex numbers  $(H_\alpha \Psi, \Psi)_{L_2(\Omega)}$  where  $\Psi$  changes over all  $\Psi \in \mathfrak{D}(H_\alpha)$  with  $\|\Psi\|_{L_2(\Omega)} = 1$ , is contained in the set  $\Xi_\alpha$ . Hence, in view of general results about numerical range (cf [18, Sec. V.3.2]), the exterior of the numerical range of  $H_\alpha$  is a connected set, and one indeed has (2.5) and (2.6).

## 4 The unperturbed waveguide

In this section we consider the case of uniform boundary conditions in the sense that  $\alpha$  is supposed to be identically equal to a constant  $\alpha_0 \in \mathbb{R}$ . We prove the spectral result (2.8) by using the fact that  $H_{\alpha_0}$  can be decomposed into a sum of the “longitudinal” operator  $-\Delta^\mathbb{R}$ , *i.e.* the self-adjoint Laplacian in  $L_2(\mathbb{R})$ , and the “transversal” operator  $-\Delta_{\alpha_0}^I$  defined in (2.10).

### 4.1 The transversal operator

We summarize here some of the results established in [20] and refer to that reference for more details.

The adjoint of  $-\Delta_{\alpha_0}^I$  is simply obtained by the replacement  $\alpha_0 \mapsto -\alpha_0$ , *i.e.*,  $(-\Delta_{\alpha_0}^I)^* = -\Delta_{-\alpha_0}^I$ . Consequently,  $(-\Delta_{\alpha_0}^I)^*$  has the same spectrum (2.11) and the corresponding set of eigenfunctions  $\{\phi_j\}_{j=0}^\infty$  can be chosen as

$$\phi_j(x_2) := \overline{A_j \psi_j(x_2)},$$

where  $\{\psi_j\}_{j=0}^\infty$  have been introduced in (2.13) and  $A_j$  are normalization constants. Choosing

$$A_{j_0} := \frac{2i\alpha_0}{1 - \exp(-2i\alpha_0 d)}, \quad A_{j_1} := \frac{2\mu_1^2}{(\mu_1^2 - \alpha_0^2)d}, \quad A_j := \frac{2\mu_j^2}{(\mu_j^2 - \alpha_0^2)d},$$

where  $j \geq 2$ ,  $(j_0, j_1) = (0, 1)$  if  $|\alpha_0| < \pi/d$  and  $(j_0, j_1) = (1, 0)$  if  $|\alpha_0| > \pi/d$  (if  $\alpha_0 = 0$ , the fraction in the definition of  $A_{j_0}$  should be understood as the expression obtained after taking the limit  $\alpha_0 \rightarrow 0$ ), we have the biorthonormality relations

$$\forall j, k \in \mathbb{N}, \quad (\psi_j, \phi_k)_{L_2(I)} = \delta_{jk}$$

together with the biorthonormal-basis-type expansion (cf [20, Prop. 4])

$$\forall \psi \in L_2(I), \quad \psi = \sum_{j=0}^{\infty} (\psi, \phi_j)_{L_2(I)} \psi_j. \quad (4.1)$$

Let us show that (4.1) can be extended to  $L_2(\Omega)$ .

**Lemma 4.1.** *For any  $\Psi \in L_2(\Omega)$ , the identity*

$$\Psi(x) = \sum_{j=0}^{\infty} \Psi_j(x_1) \psi_j(x_2) \quad \text{with} \quad \Psi_j(x_1) := (\Psi(x_1, \cdot), \phi_j)_{L_2(I)}$$

*holds true in the sense of  $L_2(\Omega)$ -norm.*

*Proof.* In view of (4.1), the series converges to  $\Psi$  in  $L_2(I)$  for almost every  $x_1 \in \mathbb{R}$ . We use the dominated convergence theorem to prove that the convergence actually holds in the norm of  $L_2(\Omega)$ . To do so, it is sufficient to check that the  $L_2(I)$ -norm of the partial sums can be uniformly estimated by a function from  $L_2(\mathbb{R})$ .

Let us introduce  $\chi_j^D(x_2) := \sin(\pi j x_2/d)$  and  $\chi_j^N(x_2) := \cos(\pi j x_2/d)$  for  $j \geq 1$ , and  $\chi_0^N(x_2) := 1/\sqrt{2}$ . We recall that  $\{\sqrt{2/d}\chi_j^D\}_{j=1}^\infty$  and  $\{\sqrt{2/d}\chi_j^N\}_{j=0}^\infty$  form complete orthonormal families in  $L_2(I)$ . Expressing  $\psi_j$  in terms of  $\chi_j^N$  and  $\chi_j^D$  for  $j \geq 2$ , and using the orthonormality, we have ( $n \geq 2$ )

$$\begin{aligned} \left\| \sum_{j=2}^n \Psi_j(x_1) \psi_j \right\|_{L_2(I)}^2 &\leq d \left\| \sum_{j=2}^n \Psi_j(x_1) \chi_j^N \right\|_{L_2(I)}^2 + d \alpha_0^2 \left\| \sum_{j=2}^n \Psi_j(x_1) \chi_j^D / \mu_j \right\|_{L_2(I)}^2 \\ &= d \sum_{j=2}^n |\Psi_j(x_1)|^2 + d \alpha_0^2 \sum_{j=2}^n |\Psi_j(x_1)|^2 / \mu_j^2 \\ &\leq d \left( 1 + \frac{\alpha_0^2}{\mu_2^2} \right) \sum_{j=2}^\infty |\Psi_j(x_1)|^2. \end{aligned} \quad (4.2)$$

Next, writing ( $j \geq 2$ )

$$\Psi_j = \sqrt{\frac{d}{2}} A_j \left( \Psi_j^N - i \frac{\alpha_0}{\mu_j} \Psi_j^D \right) \quad \text{with} \quad \Psi_j^\#(x_1) := (\Psi(x_1, \cdot), \chi_j^\#)_{L_2(I)},$$

noticing that  $|A_j| \leq c$  (valid for all  $j \geq 0$ ) where  $c$  is a constant depending uniquely on  $|\alpha_0|$  and  $d$ , and using the Parseval identities for  $\chi_j^N$  and  $\chi_j^D$ , we obtain

$$\begin{aligned} \sum_{j=2}^\infty |\Psi_j(x_1)|^2 &\leq c^2 d \sum_{j=2}^\infty \left( |\Psi_j^N|^2 + \frac{\alpha_0^2}{\mu_j^2} |\Psi_j^D|^2 \right) \\ &\leq c^2 d \left( 1 + \frac{\alpha_0^2}{\mu_2^2} \right) \|\Psi(x_1, \cdot)\|_{L_2(I)}^2. \end{aligned} \quad (4.3)$$

At the same time, using just the estimates  $|\psi_j|^2 \leq (1 + \alpha_0^2/\mu_j^2)$  valid for all  $j \geq 0$ , we readily get

$$\left\| \sum_{j=0}^1 \Psi_j(x_1) \psi_j \right\|_{L_2(I)} \leq 2 c d \left( 1 + \frac{\alpha_0^2}{\mu_0^2} \right) \|\Psi(x_1, \cdot)\|_{L_2(I)}. \quad (4.4)$$

Summing up,

$$\left\| \sum_{j=0}^n \Psi_j(x_1) \psi_j \right\|_{L_2(I)} \leq C \|\Psi(x_1, \cdot)\|_{L_2(I)} \in L_2(\mathbb{R}),$$

where  $C$  is a constant independent of  $n$ , and the usage of the dominated convergence theorem is justified.  $\square$

## 4.2 Spectrum of the unperturbed waveguide

First we show that the spectrum of  $H_{\alpha_0}$  is purely essential. Since the residual spectrum is always empty due to Corollary 2.1, it is enough to show that there are no eigenvalues.

**Proposition 4.1.** *Let  $\alpha_0 \in \mathbb{R}$  satisfy (2.12). Then*

$$\sigma_p(H_{\alpha_0}) = \emptyset.$$

*Proof.* Suppose that  $H_{\alpha_0}$  possesses an eigenvalue  $\lambda$  with eigenfunction  $\Psi$ . Multiplying the eigenvalue equation with  $\overline{\phi_j}$  and integrating over  $I$ , we arrive at the equations

$$-\Psi_j'' = (\lambda - \mu_j^2)\Psi_j \quad \text{in} \quad \mathbb{R}, \quad j \geq 0,$$

where  $\Psi_j$  are the coefficients of Lemma 4.1. Since  $\Psi_j \in L_2(\mathbb{R})$  due to Fubini's theorem, each of the equations has just a trivial solution. This together with Lemma 4.1 yields  $\Psi = 0$ , a contradiction. That is, the point spectrum of  $H_{\alpha_0}$  is empty.  $\square$

*Remark 4.1.* Regardless of the technical hypothesis (2.12), the set of isolated eigenvalues of  $H_{\alpha_0}$  is always empty. This follows from Proposition 4.1 and the fact that  $\alpha_0 \mapsto H_{\alpha_0}$  forms a holomorphic family of operators [18, Sec. VII.4].

It is well known that the spectrum of the “longitudinal” operator  $-\Delta^{\mathbb{R}}$  is also purely essential and equal to the semi-axis  $[0, +\infty)$ . In view of the separation of variables, it is reasonable to expect that the (essential) spectrum of  $H_{\alpha_0}$  will be given by that semi-axis shifted by the first eigenvalue of  $-\Delta_{\alpha_0}^I$ . First we show that the resulting interval indeed belongs to the spectrum of  $H_{\alpha_0}$ .

**Lemma 4.2.** *Let  $\alpha_0 \in \mathbb{R}$ . Then  $[\mu_0^2, +\infty) \subseteq \sigma_e(H_{\alpha_0})$ .*

*Proof.* Since the spectrum of  $H_{\alpha_0}$  is purely essential, it can be characterized by means of singular sequences [11, Thm. IX.1.3] (it is in fact an equivalent definition of another type of essential spectrum, which is in general intermediate between the essential spectra due to Wolf and Kato, and therefore coinciding with them in our case, cf Remark 2.1). That is,  $\lambda \in \sigma_e(H_{\alpha_0})$  if, and only if, there exists a sequence  $\{u_n\}_{n=1}^\infty \subset \mathfrak{D}(H_{\alpha_0})$  such that  $\|u_n\|_{L_2(\Omega)} = 1$ ,  $u_n \rightharpoonup 0$  and  $\|H_{\alpha_0}u_n - \lambda u_n\|_{L_2(\Omega)} \rightarrow 0$  as  $n \rightarrow \infty$ . Let  $\{\varphi_n\}_{n=1}^\infty$  be a singular sequence of  $-\Delta^{\mathbb{R}}$  corresponding to a given  $z \in [0, +\infty)$ . Then it is easy to verify that  $u_n$  defined by  $u_n(x) := \varphi_n(x_1)\psi_0(x_2)/\|\psi_0\|_{L_2(I)}$  forms a singular sequence of  $H_{\alpha_0}$  corresponding to  $z + \mu_0^2$ .  $\square$

To get the opposite inclusion, we employ the fact that the biorthonormal-basis-type relations (4.1) are available. This enables us to decompose the resolvent of  $H_{\alpha_0}$  into the transverse biorthonormal-basis.

**Lemma 4.3.** *Let  $\alpha_0 \in \mathbb{R}$  satisfy (2.12). Then  $\mathbb{C} \setminus [\mu_0^2, +\infty) \subseteq \rho(H_{\alpha_0})$  and for any  $z \in \mathbb{C} \setminus [\mu_0^2, +\infty)$  we have*

$$(H_{\alpha_0} - z)^{-1} = \sum_{j=0}^{\infty} (-\Delta^{\mathbb{R}} + \mu_j^2 - z)^{-1} B_j.$$

Here  $B_j$  is a bounded operator on  $L_2(\Omega)$  defined by

$$(B_j \Psi)(x) := (\Psi(x_1, \cdot), \phi_j)_{L_2(I)} \psi_j(x_2), \quad \Psi \in L_2(\Omega),$$

and  $(-\Delta^{\mathbb{R}} + \mu_j^2 - z)^{-1}$  abbreviates  $(-\Delta^{\mathbb{R}} + \mu_j^2 - z)^{-1} \otimes 1$  on  $L_2(\mathbb{R}) \otimes L_2(I) \simeq L_2(\Omega)$ .

*Proof.* Put  $z \in \mathbb{C} \setminus [\mu_0^2, +\infty)$ . For every  $\Psi \in L_2(\Omega)$  and all  $j \geq 0$  we denote  $U_j := (-\Delta^{\mathbb{R}} + \mu_j^2 - z)^{-1} \Psi_j \in L_2(\mathbb{R})$ , where  $\Psi_j$  are defined in Lemma 4.1. It is clear that

$$\|U_j\|_{L_2(\mathbb{R})} \leq \frac{\|\Psi_j\|_{L_2(\mathbb{R})}}{\text{dist}(z, [\mu_j^2, +\infty))} \leq C \frac{\|\Psi_j\|_{L_2(\mathbb{R})}}{j^2 + 1}, \quad \|U_j'\|_{L_2(\mathbb{R})} \leq C \frac{\|\Psi_j\|_{L_2(\mathbb{R})}}{\sqrt{j^2 + 1}}, \quad (4.5)$$

where  $C$  is a constant depending uniquely on  $|\alpha_0|$ ,  $d$  and  $z$ ; the second inequality follows from the identity  $\|U_j'\|_{L_2(\mathbb{R})}^2 + (\mu_j^2 - z)\|U_j\|_{L_2(\mathbb{R})}^2 = (\Psi_j, U_j)_{L_2(\mathbb{R})}$ . Using (4.5) and estimates of the type (4.4), it is readily seen that each function  $R_j : x \mapsto U_j(x_1)\psi_j(x_2)$  belongs to  $W_2^1(\Omega)$ . We will show that it is the case for their infinite sum too. Firstly, a consecutive use of (4.2), the first inequality of (4.5) and (4.3) together with Fubini's theorem implies  $\|\sum_{j=2}^n R_j\|_{L_2(\Omega)} \leq K \|\Psi\|_{L_2(\Omega)}$ , where  $K$  is a constant independent of  $n \geq 2$ . Secondly, a similar estimate for the partial sum of  $\partial_1 R_j$  can be obtained in the same way, provided that one uses the second inequality of (4.5) now. Finally, since the derivative of  $\psi_j$  as well can be expressed in terms of  $\chi_j^N$  and  $\chi_j^D$  introduced in the proof of Lemma 4.1, a consecutive use of the estimates of the type (4.2) and the first inequality of (4.5) together with Fubini's theorem implies

$$\left\| \sum_{j=2}^n \partial_2 R_j \right\|_{L_2(\Omega)}^2 \leq d \sum_{j=2}^n (\alpha_0^2 + \mu_j^2) \|U_j\|_{L_2(\mathbb{R})}^2 \leq d C^2 \sum_{j=2}^n \frac{\alpha_0^2 + \mu_j^2}{(j^2 + 1)^2} \|\Psi_j\|_{L_2(\mathbb{R})}^2;$$

here the fraction in the upper bound forms a bounded sequence in  $j$ , so that we may continue to estimate as above using (4.3) together with Fubini's theorem again. Summing up, the series  $\sum_{j=0}^{\infty} R_j$  converges in  $W_2^1(\Omega)$  to a function  $R$  and

$$\|R\|_{W_2^1(\Omega)} \leq \tilde{K} \|\Psi\|_{L_2(\Omega)},$$

where  $\tilde{K}$  depends uniquely on  $|\alpha_0|$ ,  $d$  and  $z$ . Employing this fact and the definition of  $U_j$ , one can check easily that  $R$  satisfies the identity  $h_{\alpha_0}(R, \Phi) - z(R, \Phi)_{L_2(\Omega)} = (\Psi, \Phi)_{L_2(\Omega)}$  for all  $\Phi \in W_2^1(\Omega)$ . It implies that  $R \in \mathfrak{D}(H_{\alpha_0})$  and  $(H_{\alpha_0} - z)R = \Psi$ , i.e.,  $R = (H_{\alpha_0} - z)^{-1} \Psi$ .  $\square$

Lemmata 4.2 and 4.3 yield

**Proposition 4.2.** *Let  $\alpha_0 \in \mathbb{R}$ . Then*

$$\sigma_e(H_{\alpha_0}) = [\mu_0^2, +\infty).$$

*Proof.* In view of the lemmata, the result holds for every  $\alpha_0 \in \mathbb{R}$  except for a discrete set of points complementary to the hypothesis (2.12). However, these points can be included by noticing that  $\alpha_0 \mapsto H_{\alpha_0}$  forms a holomorphic family of operators (cf Remark 4.1).  $\square$

Proposition 4.2 and Proposition 4.1 together with Remark 4.1 imply that the spectrum of  $H_{\alpha_0}$  is real and (2.8) holds true for every  $\alpha_0 \in \mathbb{R}$ .

*Remark 4.2.* It follows from (2.8) that the spectrum of  $H_{\alpha_0}$  is equal to the sum of the spectra of  $-\Delta^{\mathbb{R}}$  and  $-\Delta_{\alpha_0}^I$ . This result could alternatively be obtained by using a general theorem about the spectrum of tensor products [25, Thm. XIII.35] and the fact that the one-dimensional operators generate bounded holomorphic semigroups. However, we do not use this way of proof since Lemma 4.3 is employed not only in the proof of (2.8) but also in the proofs of Theorems 2.5 and 2.6.

## 5 Stability of the essential spectrum

In this section we show that the essential spectrum is stable under a compactly supported perturbation of the boundary conditions. In fact, we will establish a stronger result, namely that the difference of the resolvents of  $H_\alpha$  and  $H_{\alpha_0}$  is a compact operator. As an auxiliary result, we shall need the following lemma.

**Lemma 5.1.** *Let  $\alpha_0 \in \mathbb{R}$  and  $\varphi \in L_2(\partial\Omega)$ . There exist positive constants  $c$  and  $C$ , depending on  $d$  and  $|\alpha_0|$ , such that any solution  $\Psi \in W_2^1(\Omega)$  of the boundary value problem*

$$\begin{cases} (-\Delta - z)\Psi = 0 & \text{in } \Omega, \\ (\partial_2 + i\alpha_0)\Psi = \varphi & \text{on } \partial\Omega, \end{cases} \quad (5.1)$$

with any  $z \leq -c$ , satisfies the estimate

$$\|\Psi\|_{W_2^1(\Omega)} \leq C \|\varphi\|_{L_2(\partial\Omega)}. \quad (5.2)$$

*Proof.* Multiplying the first equation of (5.1) by  $\overline{\Psi}$  and integrating over  $\Omega$ , we arrive at the identity

$$\|\nabla\Psi\|^2 - z\|\Psi\|^2 + i\alpha_0 \int_{\partial\Omega} \nu_2 |\Psi|^2 - \int_{\partial\Omega} \nu_2 \varphi \overline{\Psi} = 0, \quad (5.3)$$

where  $\nu_2$  denotes the second component of the outward unit normal vector to  $\partial\Omega$ . Using the Schwarz and Cauchy inequalities together with  $|\nu_2| = 1$ , we have

$$\begin{aligned} \left| \int_{\partial\Omega} \nu_2 |\Psi|^2 \right| &= \left| \int_{\partial\Omega} \partial_2 |\Psi|^2 \right| = 2 |\operatorname{Re}(\Psi, \partial_2 \Psi)| \leq \delta^{-1} \|\Psi\|^2 + \delta \|\nabla\Psi\|^2, \\ 2 \left| \int_{\partial\Omega} \nu_2 \varphi \overline{\Psi} \right| &\leq \delta^{-1} \|\varphi\|_{L_2(\partial\Omega)}^2 + \delta \|\Psi\|_{L_2(\partial\Omega)}^2, \end{aligned}$$

with any  $\delta \in (0, 1)$ . Here  $\|\Psi\|_{L_2(\partial\Omega)} \leq C \|\Psi\|_{W_2^1(\Omega)}$ , where  $C$  is the constant coming from the embedding of  $W_2^1(\Omega)$  in  $L_2(\partial\Omega)$  (depending only on  $d$  in our case). Choosing now sufficiently small  $\delta$  and sufficiently large negative  $z$ , it is clear that (5.3) can be cast into the inequality (5.2).  $\square$

Now we are in a position to prove

**Proposition 5.1.** *Let  $\alpha - \alpha_0 \in C_0(\mathbb{R}) \cap W_\infty^1(\mathbb{R})$  with  $\alpha_0 \in \mathbb{R}$ . Then*

$$(H_\alpha - z)^{-1} - (H_{\alpha_0} - z)^{-1} \text{ is compact in } L_2(\Omega)$$

for any  $z \in \rho(H_\alpha) \cap \rho(H_{\alpha_0})$ .

*Proof.* It is enough to prove the result for one  $z$  in the intersection of the resolvent sets of  $H_\alpha$  and  $H_{\alpha_0}$ , and we can assume that the one is negative (since the operators are  $m$ -accretive). Given  $\Phi \in L_2(\Omega)$ , let  $\Psi := (H_\alpha - z)^{-1}\Phi - (H_{\alpha_0} - z)^{-1}\Phi$ . It is easy to check that  $\Psi$  is the unique solution to (5.1) with  $\varphi := -i(\alpha - \alpha_0)T(H_\alpha - z)^{-1}\Phi$ , where  $T$  denotes the trace operator from  $W_2^2(\Omega) \supset \mathfrak{D}(H_\alpha)$  to  $W_2^1(\partial\Omega)$ . By virtue of Lemma 5.1, it is therefore enough to show that  $(\alpha - \alpha_0)T(H_\alpha - z)^{-1}$  is a compact operator from  $L_2(\Omega)$  to  $L_2(\partial\Omega)$ . However, this property follows from the fact that  $W_2^1(\partial\Omega)$  is compactly embedded in  $L_2(\omega)$  for every bounded subset  $\omega$  of  $\partial\Omega$ , due to the Rellich-Kondrachov theorem [1, Sec. VI].  $\square$

**Corollary 5.2.** *Suppose the hypothesis of Proposition 5.1. Then*

$$\sigma_e(H_\alpha) = [\mu_0^2, +\infty).$$

*Proof.* Our definition of essential spectrum is indeed stable under relatively compact perturbations [11, Thm. IX.2.4].  $\square$

## 6 Point spectrum

In this section we prove Theorems 2.3–2.6 and Propositions 2.1–2.2. In the proofs of Theorems 2.4–2.6 we follow the main ideas of [13]. Throughout this section we assume that the identity (2.14) and the assumption (2.12) hold true.

### 6.1 Proof of Theorem 2.3

Any eigenvalue of infinite geometric multiplicity belongs to the essential spectrum which is real by Theorem 2.2. Let  $\lambda$  be an eigenvalue of  $H_\alpha$  of finite geometric multiplicity, and  $\Psi$  be an associated eigenfunction. Using that  $\alpha$  is of compact support, it is easy to check that  $x \mapsto \Psi(-x_1, d - x_2)$  is an eigenfunction associated with  $\lambda$ , too. The geometric multiplicity of  $\lambda$  being finite, we conclude that at least one of the eigenfunction associated with  $\lambda$  satisfies  $|\Psi(x)| = |\Psi(-x_1, d - x_2)|$ . Taking this identity into account, integrating by parts and using the hypothesis that  $\alpha$  is odd, we obtain

$$\lambda \|\Psi\|_{L_2(\Omega)}^2 = h_\alpha^1[\Psi] + i h_\alpha^2[\Psi] = \|\nabla \Psi\|_{L_2(\Omega)}^2,$$

which implies that  $\lambda$  is real.  $\square$

### 6.2 Auxiliary results

Let a function  $F \in L_2(\Omega)$  be such that  $\text{supp } F \subseteq \overline{\Omega_b}$  for fixed  $b > 0$ . We consider the boundary value problem

$$\begin{cases} -\Delta U = (\mu_0^2 - k^2)U + F & \text{in } \Omega, \\ (\partial_2 + i\alpha_0)U = 0 & \text{on } \partial\Omega, \end{cases} \quad (6.1)$$

where the parameter  $k \in \mathbb{C}$  ranges in a small neighbourhood of zero. The problem can be solved by separation of variables justified in Lemma 4.3 whenever  $k^2 \notin (-\infty, 0]$ . Moreover, it is possible to extend the solution of (6.1) analytically with respect to  $k$ . Namely, the following statement is valid.

**Lemma 6.1.** *For all small  $k \in \mathbb{C}$  there exists the unique solution to (6.1) satisfying*

$$U(x; k) = c_\pm(k) e^{-k|x_1|} \psi_0(x_2) + \mathcal{O}(e^{-\text{Re } \sqrt{\mu_1^2 - \mu_0^2 + k^2}|x_1|}), \quad (6.2)$$

*in the limit  $x_1 \rightarrow \pm\infty$ , where  $c_\pm(k)$  are constants. The mapping  $T_1(k)$  defined as  $T_1(k)F := U$  is a bounded linear operator from  $L_2(\Omega_b)$  into  $W_2^2(\Omega_a)$  for each  $a > 0$ . This operator is meromorphic with respect to  $k$  and has the simple pole at zero,*

$$T_1(k)F = \frac{\psi_0}{2k} \int_{\Omega} F(x) \overline{\phi_0}(x_2) dx + T_2(k)F,$$

where for each  $a > 0$  the operator  $T_2(k) : L_2(\Omega_b) \rightarrow W_2^2(\Omega_a)$  is linear, bounded and holomorphic with respect to  $k$  small enough. The function  $\widehat{U} := T_2(0)F$  is the unique solution to the problem

$$\begin{cases} -\Delta \widehat{U} = \mu_0^2 \widehat{U} + F & \text{in } \Omega, & (\partial_2 + i\alpha_0) \widehat{U} = 0 & \text{on } \partial\Omega, \\ \widehat{U}(x) = -\frac{\psi_0(x_2)}{2} \int_{\Omega} |x_1 - t_1| F(t) \overline{\phi_0}(t_2) dt + \mathcal{O}(e^{-\sqrt{\mu_1^2 - \mu_0^2}|x_1|}), & |x_1| \rightarrow +\infty, \end{cases} \quad (6.3)$$

given by the formula

$$\widehat{U}(x) = \sum_{j=0}^{\infty} \widehat{U}_j(x_1) \psi_j(x_2) \quad (6.4)$$

with

$$\widehat{U}_j(x_1) := \begin{cases} -\frac{1}{2} \int_{\Omega} |x_1 - t_1| F(t) \overline{\phi_0}(t_2) dt & \text{if } j = 0, \\ \frac{1}{2\sqrt{\mu_j^2 - \mu_0^2}} \int_{\Omega} e^{-\sqrt{\mu_j^2 - \mu_0^2}|x_1 - t_1|} F(t) \overline{\phi_j}(t_2) dt & \text{if } j \geq 1. \end{cases}$$

The lemma is proved in the same way as Lemma 3.1 in [4].

Let  $M_\varepsilon$  be the operator of multiplication by the function  $x \mapsto e^{-i\varepsilon\beta(x_1)x_2}$ . It is straightforward to check that  $H_\alpha$  is unitarily equivalent to the operator

$$M_\varepsilon^{-1} H_\alpha M_\varepsilon = H_{\alpha_0} - \varepsilon L_\varepsilon,$$

where

$$L_\varepsilon = -2i\beta'(x_1)x_2 \frac{\partial}{\partial x_1} - 2i\beta(x_1) \frac{\partial}{\partial x_2} - (\varepsilon\beta^2(x_1) + i\beta''(x_1)x_2 + \varepsilon\beta'^2(x_1)x_2^2).$$

We observe that the coefficients of  $L_\varepsilon$  are compactly supported and that their supports are bounded uniformly in  $\varepsilon$ .

It follows that the eigenvalue equation for  $H_\alpha$  is equivalent to

$$H_{\alpha_0} U = \lambda U + \varepsilon L_\varepsilon U,$$

where an eigenfunction  $\Psi$  of  $H_\alpha$  satisfies  $\Psi = M_\varepsilon U$ . It can be rewritten as

$$\begin{cases} -\Delta U = (\mu_0^2 - k^2)U + \varepsilon L_\varepsilon U & \text{in } \Omega, \\ (\partial_2 + i\alpha_0)U = 0 & \text{on } \partial\Omega, \end{cases} \quad (6.5)$$

where we have replaced  $\lambda$  by  $\mu_0^2 - k^2$ .

Now, let  $\lambda$  be an eigenvalue for  $H_\alpha$  close to  $\mu_0^2$ . As  $x_1 \rightarrow \pm\infty$ , the solution  $U$  to (6.5) satisfies the asymptotic formula (6.2), where  $k = \sqrt{\mu_0^2 - \lambda}$  and the branch of the root is specified by the requirement  $\operatorname{Re} k > 0$ . Such restriction guarantees that the function  $U$  together with their derivatives decays exponentially at infinity and thus belongs to  $W_2^2(\Omega)$ . Hence, the set of all  $k$  for which the problem (6.5), (6.2) has a nontrivial solution includes the set of all values of  $k$  related to the eigenvalues of  $H_\alpha$  by the relation  $\lambda = \mu_0^2 - k^2$ . Thus, it is sufficient to find all small  $k$  for which a nontrivial solution to (6.5), (6.2) exists and

to check whether the solution belongs to  $W_2^2(\Omega)$ . If it does, the corresponding number  $\lambda = \mu_0^2 - k^2$  is an eigenvalue of  $H_\alpha$ .

We introduce the numbers

$$k_1(\varepsilon) := \frac{1}{2} \int_{\Omega} \overline{\phi_0}(x_2) (L_\varepsilon \psi_0)(x) \, dx, \quad k_2(\varepsilon) := \frac{1}{2} \int_{\Omega} \overline{\phi_0}(x_2) (L_\varepsilon T_2(0) L_\varepsilon \psi_0)(x) \, dx.$$

Basing on Lemma 6.1 and arguing in the same way as in [13, Sec. 2] one can prove easily the following statement (see also [4, Sec. 4]).

**Lemma 6.2.** *There exists the unique function  $\varepsilon \mapsto k(\varepsilon)$  converging to zero as  $\varepsilon \rightarrow +0$  for which the problem (6.5), (6.2) has a nontrivial solution. It satisfies the asymptotics*

$$k(\varepsilon) = \varepsilon k_1(\varepsilon) + \varepsilon^2 k_2(\varepsilon) + \mathcal{O}(\varepsilon^3).$$

*The associated nontrivial solution to (6.5), (6.2) is unique up to a multiplicative constant and can be chosen so that it obeys (6.2) with*

$$c_\pm(k(\varepsilon)) = 1 + \mathcal{O}(\varepsilon), \quad \varepsilon \rightarrow +0, \quad (6.6)$$

as well as

$$U(x; \varepsilon) = \psi_0(x_2) + \mathcal{O}(\varepsilon)$$

in  $W_2^2(\Omega_a)$  for each fixed  $a > 0$ .

### 6.3 Proof of Theorems 2.5 and 2.6

It follows from Lemma 6.2 that there is at most one simple eigenvalue of  $H_\alpha$  converging to  $\mu_0^2$  as  $\varepsilon \rightarrow +0$ . A sufficient condition guaranteeing the existence of such eigenvalue is the inequality

$$\operatorname{Re}(k_1(\varepsilon) + \varepsilon k_2(\varepsilon)) \geq C(\varepsilon) \varepsilon^2, \quad C(\varepsilon) \rightarrow +\infty, \quad \varepsilon \rightarrow +0, \quad (6.7)$$

that is implied by (6.2), (6.6), the definition of the operator  $M_\varepsilon$  and the assumption on  $\beta$ . The sufficient condition of the absence of the eigenvalue is the opposite inequality

$$\operatorname{Re}(k_1(\varepsilon) + \varepsilon k_2(\varepsilon)) \leq -C(\varepsilon) \varepsilon^2, \quad C(\varepsilon) \rightarrow +\infty, \quad \varepsilon \rightarrow +0. \quad (6.8)$$

Thus, we just need to calculate the numbers  $k_1$  and  $k_2$  to prove the theorems.

It is easy to compute the coefficient  $k_1$ ,

$$k_1(\varepsilon) = \begin{cases} -\alpha_0 \langle \beta \rangle + k_1'(0) \varepsilon & \text{if } |\alpha_0| < \pi/d, \\ k_1'(0) \varepsilon & \text{if } |\alpha_0| > \pi/d, \end{cases} \quad (6.9)$$

where

$$k_1'(0) := -\frac{1}{2} \int_{\Omega} \psi_0(x_2) \overline{\phi_0}(x_2) (\beta^2(x_1) + \beta'^2(x_1) x_2^2) \, dx.$$

It is more complicated technically to calculate  $k_2$ . This coefficient depends on  $\varepsilon$  as well; to prove the theorem we need the leading term of its asymptotics as  $\varepsilon \rightarrow +0$ . We begin the calculations by observing an obvious identity,

$$L_\varepsilon U = L_0 U + \mathcal{O}(\varepsilon),$$

where

$$\begin{aligned}(L_0 U)(x) &:= -2i\beta'(x_1)x_2 \frac{\partial U(x)}{\partial x_1} - 2i\beta(x_1) \frac{\partial U(x)}{\partial x_2} - i\beta''(x_1)x_2 U(x) \\ &= i\left(\beta(x_1)x_2 \Delta U(x) - \Delta[\beta(x_1)x_2 U(x)]\right),\end{aligned}$$

which is valid for each  $U \in W_{2,loc}^2(\Omega)$  in  $L_2(\Omega_a)$ , if  $a$  is large enough and independent of  $\varepsilon$ . Thus,

$$k_2(\varepsilon) = k_2(0) + \mathcal{O}(\varepsilon), \quad \text{where} \quad k_2(0) = \frac{1}{2} \int_{\Omega} \overline{\phi_0}(x_2) (L_0 T_2(0) L_0 \psi_0)(x) dx.$$

We denote  $\widehat{U} := T_2(0)\widehat{F}$  and  $\widehat{F} := L_0 \psi_0$ . Taking into account the problem (6.3) for  $\widehat{U}$  and integrating by parts, we obtain

$$\begin{aligned}k_2(0) &= \frac{i}{2} \int_{\Omega} \overline{\phi_0}(x_2) \left( \beta(x_1)x_2 \Delta \widehat{U}(x) - \Delta[\beta(x_1)x_2 \widehat{U}(x)] \right) dx \\ &= -\frac{i}{2} \int_{\Omega} \overline{\phi_0}(x_2) (\Delta + \mu_0^2) \beta(x_1)x_2 \widehat{U}(x) dx - \frac{i}{2} \int_{\Omega} \overline{\phi_0}(x_2) \beta(x_1)x_2 \widehat{F}(x) dx \\ &= -\frac{i}{2} \left\langle \beta \left[ \overline{\phi_0}(d) \widehat{U}(\cdot, d) - \overline{\phi_0}(0) \widehat{U}(\cdot, 0) \right] \right\rangle - \frac{i}{2} \int_{\Omega} \overline{\phi_0}(x_2) \beta(x_1)x_2 \widehat{F}(x) dx.\end{aligned}\tag{6.10}$$

The last term on the right hand side of this identity is calculated by integration by parts,

$$\begin{aligned}-\frac{i}{2} \int_{\Omega} \overline{\phi_0}(x_2) \beta(x_1)x_2 \widehat{F}(x) dx &= -\frac{1}{2} \int_{\Omega} \overline{\phi_0}(x_2) \beta(x_1)x_2 [2\beta(x_1)\psi_0'(x_2) + \beta''(x_1)x_2\psi_0(x_2)] dx \\ &= -\frac{1}{2} \int_{\Omega} \beta^2(x_1)x_2 (\overline{\phi_0}\psi_0)'(x_2) dx + \frac{1}{2} \int_{\Omega} x_2^2 \beta'^2(x_1) \overline{\phi_0}(x_2) \psi_0(x_2) dx \\ &= -\frac{\overline{\phi_0}(d)\psi_0(d)d}{2} \langle \beta^2 \rangle + \frac{1}{2} \int_{\Omega} \overline{\phi_0}(x_2) \psi_0(x_2) [\beta^2(x_1) + \beta'^2(x_1)x_2^2] dx.\end{aligned}$$

This formula, (6.9) and (6.10) yield

$$k_1(\varepsilon) + \varepsilon k_2(\varepsilon) = \begin{cases} -\varepsilon \alpha_0 \langle \beta \rangle + \varepsilon^2 K + \mathcal{O}(\varepsilon^3) & \text{if } |\alpha_0| < \pi/d, \\ \varepsilon^2 K + \mathcal{O}(\varepsilon^3) & \text{if } |\alpha_0| > \pi/d, \end{cases} \tag{6.11}$$

where

$$K := -\frac{i}{2} \left\langle \beta \left[ \overline{\phi_0}(d) \widehat{U}(\cdot, d) - \overline{\phi_0}(0) \widehat{U}(\cdot, 0) \right] \right\rangle - \frac{\overline{\phi_0}(d)\psi_0(d)d}{2} \langle \beta^2 \rangle.$$

Thus, it remains to calculate  $K$ . In order to do it, we construct the function  $\widehat{U}$  as the series (6.4).

**Case  $|\alpha_0| < \pi/d$ :** Using the identity

$$\widehat{F}(x) = -2i\beta(x_1)\psi'_0(x_2) - i\beta''(x_1)x_2\psi_0(x_2),$$

one can check that

$$\begin{aligned}\widehat{F}(x) &= -2\alpha_0\beta(x_1)\psi_0(x_2) - i\beta''(x_1)\sum_{j=0}^{\infty}c_j\psi_j(x_2), \\ \widehat{U}_j(x_1) &= \begin{cases} ic_0\beta(x_1) - 2\alpha_0v_0(x_1) & \text{if } j = 0, \\ ic_j[\beta(x_1) - (\mu_j^2 - \mu_0^2)v_j(x_1)] & \text{if } j \geq 1, \end{cases}\end{aligned}$$

where  $c_j := \int_I x_2\psi_0(x_2)\overline{\phi_j}(x_2)dx_2$  and the functions  $v_j$  were introduced in (2.15). Substituting now the formulae for  $\widehat{U}_j$  and (6.4) into (6.11), we arrive at the following chain of identities

$$\begin{aligned}K &= \frac{1}{2}\left[\sum_{j=0}^{\infty}c_j\psi_j(x_2)\overline{\phi_0}(x_2)\right]_{x_2=0}^{x_2=d}\langle\beta^2\rangle + 2\alpha_0^2\langle\beta v_0\rangle \\ &\quad - \frac{2\alpha_0}{d}\sum_{j=1}^{\infty}\frac{i[e^{i\alpha_0d} - (-1)^j]\mu_j^2}{[e^{i\alpha_0d} + (-1)^j](\mu_j^2 - \mu_0^2)}\langle\beta v_j\rangle - \frac{\overline{\phi_0}(d)\psi_0(d)d}{2}\langle\beta^2\rangle \\ &= \frac{1}{2}\left[x_2\psi_0(x_2)\overline{\phi_0}(x_2)\right]_{x_2=0}^{x_2=d}\langle\beta^2\rangle + 2\alpha_0^2\langle\beta v_0\rangle \\ &\quad + \frac{2\alpha_0}{d}\sum_{j=1}^{\infty}\frac{\mu_j^2\langle\beta v_j\rangle}{\mu_j^2 - \mu_0^2}\tan\frac{\alpha_0d + \pi j}{2} - \frac{\overline{\phi_0}(d)\psi_0(d)d}{2}\langle\beta^2\rangle,\end{aligned}$$

where the last expression coincides with  $\tau$  for  $|\alpha_0| < \pi/d$ .

**Case  $|\alpha_0| > \pi/d$ :** Following the same scheme as above, we arrive at

$$\begin{aligned}\widehat{F}(x) &= -i\beta''(x_1)\sum_{j=0}^{\infty}c_j\psi_j(x_2) - \frac{4\alpha_0\psi_1(x_2)\beta(x_1)}{1 - e^{-i\alpha_0d}} \\ &\quad - \frac{2i}{d}(\mu_0^2 - \mu_1^2)\sum_{j=1}^{\infty}\frac{\mu_{2j}^2\psi_{2j}(x_2)}{(\mu_{2j}^2 - \mu_1^2)(\mu_{2j}^2 - \mu_0^2)}\beta(x_1), \\ \widehat{U}_j(x_1) &= \begin{cases} ic_0\beta(x_1) & \text{if } j = 0, \\ ic_1\beta(x_1) - \frac{2\alpha_0}{1 - e^{-i\alpha_0d}}v_1(x_1) & \text{if } j = 1, \\ ic_j\beta(x_1) + \frac{2i\mu_j^2[1 + (-1)^j]}{(\mu_j^2 - \mu_1^2)d}v_j(x_j) & \text{if } j \geq 2, \end{cases}\end{aligned}$$

and check that  $K = \tau$  for  $|\alpha_0| > \pi/d$ .

The series in the formulae for  $\tau$  converge since the functions  $v_j$  satisfy

$$-v_j'' + (\mu_j^2 - \mu_0^2)v_j = \beta \quad \text{in } \mathbb{R}, \quad (6.12)$$

and by [18, Ch. V, §3.5, Formula (3.16)] ( $j \geq 1$ )

$$|\langle\beta v_j\rangle| \leq \|\beta\|_{L_2(\mathbb{R})}\|v\|_{L_2(\mathbb{R})} \leq \frac{\|\beta\|_{L_2(\mathbb{R})}^2}{\mu_j^2 - \mu_0^2}. \quad (6.13)$$

Summing up,

$$k_1(\varepsilon) + \varepsilon k_2(\varepsilon) = \begin{cases} -\varepsilon \alpha_0 \langle \beta \rangle + \varepsilon^2 \tau + \mathcal{O}(\varepsilon^3) & \text{if } |\alpha_0| < \pi/d, \\ \varepsilon^2 \tau + \mathcal{O}(\varepsilon^3) & \text{if } |\alpha_0| > \pi/d. \end{cases}$$

All the statements of the theorems – except for the reality of the eigenvalue – follow from these formulae, the identity  $\lambda = \mu_0^2 - k^2$ , the inequalities (6.7) and (6.8), Lemma 6.2, the asymptotics (6.2) for  $U$ , and the definition of the operator  $M_\varepsilon$ .

Let us show that  $\lambda_\varepsilon$  is necessarily real as  $\varepsilon \rightarrow 0+$ . Let  $\Psi_\varepsilon$  be the eigenfunction associated with the eigenvalue  $\lambda_\varepsilon$ . It is easy to check that the function  $x \mapsto \bar{\Psi}_\varepsilon(x_1, d - x_2)$  is an eigenfunction of  $H_\alpha$  associated with the eigenvalue  $\bar{\lambda}_\varepsilon$ . This eigenvalue converges to  $\mu_0^2$  as  $\varepsilon \rightarrow +0$ . By the uniqueness of such eigenvalue we obtain  $\lambda_\varepsilon = \bar{\lambda}_\varepsilon$  that completes the proof.  $\square$

## 6.4 Proof of Theorem 2.4

We employ here the same argument as in the previous proof. The formula for  $k(\varepsilon)$  in the case  $\alpha_0 = 0$  can be obtained from that for  $|\alpha_0| < \pi/d$  by passing to the limit  $\alpha_0 \rightarrow 0$ . It leads us to the relation

$$k(\varepsilon) = \varepsilon^2 \tau + \mathcal{O}(\varepsilon^3) \quad \text{with} \quad \tau = - \sum_{j=0}^{\infty} \frac{4 \langle \beta v_{2j+1} \rangle}{\mu_{2j+1} d^2}.$$

To prove the theorem it is sufficient to show that  $\tau < 0$ . Indeed, the equation (6.12) implies that for  $j \geq 1$

$$\langle \beta v_j \rangle = \|v'_j\|_{L_2(\mathbb{R})}^2 + (\mu_j^2 - \mu_0^2) \|v_j\|_{L_2(\mathbb{R})}^2 > 0. \quad (6.14)$$

Thus,  $\tau < 0$ .  $\square$

## 6.5 Proof of Proposition 2.1

Since  $\langle \beta \rangle = 0$ , the function  $v_0$  is constant at infinity. Hence, by the equation (6.12) for  $v_0$ ,

$$\langle \beta v_0 \rangle = \|v'_0\|_{L_2(\mathbb{R})}^2.$$

At the same time, it follows from (6.13) and (6.14) that for  $j \geq 1$ ,

$$0 < \langle \beta v_j \rangle < \frac{\|\beta\|_{L_2(\mathbb{R})}^2}{\mu_j^2 - \mu_0^2}.$$

The relations obtained allow us to estimate

$$\begin{aligned} \tau &> 2\alpha_0^2 \|v'_0\|_{L_2(\mathbb{R})}^2 - \frac{2\alpha_0 \cot \frac{\alpha_0 d}{2}}{d} \|\beta\|_{L_2(\mathbb{R})}^2 \left( \frac{\mu_1^2}{(\mu_1^2 - \mu_0^2)^2} + \sum_{j=1}^{\infty} \frac{\mu_{2j+1}^2}{(\mu_{2j+1}^2 - \mu_1^2)^2} \right) \\ &= \frac{\alpha_0^2 l^3}{2} \left\| \int_{\mathbb{R}} \text{sign}(\cdot - t_1) \tilde{\beta}(t_1) dt_1 \right\|_{L_2(\mathbb{R})}^2 \\ &\quad - \frac{2\alpha_0 l \cot \frac{\alpha_0 d}{2}}{d} \left( \frac{\mu_1^2}{(\mu_1^2 - \mu_0^2)^2} + \frac{d^2}{16\pi^2} + \frac{d^2}{48} \right) \|\tilde{\beta}\|_{L_2(\mathbb{R})}^2, \end{aligned}$$

where the resulting expression is positive under the hypothesis.  $\square$

## 6.6 Proof of Proposition 2.2

Using (6.14), we obtain

$$\tau > \frac{2\alpha_0\pi^2 \cot \frac{\alpha_0 d}{2}}{(\mu_1^2 - \mu_0^2)d^3} \langle \beta v_1 \rangle + \frac{8\pi^2}{(\mu_1^2 - \mu_0^2)d^4} \sum_{j=1}^m \frac{\mu_{2j}^2 \langle \beta v_{2j} \rangle}{\mu_{2j}^2 - \mu_1^2},$$

where the right hand side is non-negative under the hypothesis (2.19).  $\square$

Let us show that the inequality (2.19) can be achieved if  $\alpha_0 \rightarrow \mu_2 + 0$ . In this case  $m = 1$ , and it is sufficient to check that

$$\alpha_0 \langle \beta v_1 \rangle \cot \frac{\alpha_0 d}{2} > \frac{16\pi^2 \langle \beta v_2 \rangle}{(\mu_1^2 - \mu_2^2)d^3} = \frac{16\pi^2 \langle \beta v_2 \rangle}{(\alpha_0^2 - \mu_2^2)d^3}. \quad (6.15)$$

It follows from the definition of  $v_1$  that it satisfies the asymptotic formula

$$v_1(x_1) = v_2(x_1) + (\mu_2 - \alpha_0) \widehat{v}(x_1) + \mathcal{O}((\mu_2 - \alpha_0)^2)$$

in  $L_2(\mathbb{R})$ -norm, where the function  $\widehat{v}$  is given by

$$\widehat{v}(x_1) := \int_{\mathbb{R}} \frac{(\sqrt{3}\mu_0|x_1 - t_1| + 1)e^{-\sqrt{3}\mu_0|x_1 - t_1|}}{3\sqrt{3}\mu_0^2} \beta(t_1) dt_1,$$

and satisfies the equation

$$-\widehat{v}'' + 3\mu_0^2 \widehat{v} = 4\mu_0 v_2 \quad \text{in } \mathbb{R}.$$

We multiply this equation by  $v_2$  and integrate by parts over  $\mathbb{R}$  taking into account the equation (6.12) for  $v_2$ ,

$$4\mu_0 \|v_2\|_{L_2(\mathbb{R})}^2 = \langle \beta \widehat{v} \rangle.$$

Hence,

$$\langle \beta v_1 \rangle = \langle \beta v_2 \rangle + 4(\mu_2 - \alpha_0)\mu_0 \|v_2\|_{L_2(\mathbb{R})}^2 + \mathcal{O}((\mu_2 - \alpha_0)^2).$$

Employing this identity, we write the asymptotic expansions for the both sides of (6.15) as  $\alpha_0 \rightarrow \mu_2 + 0$ , and obtain

$$\begin{aligned} \alpha_0 \langle \beta v_1 \rangle \cot \frac{\alpha_0 d}{2} &= \frac{4\pi \langle \beta v_2 \rangle}{(\alpha_0 - \mu_2)d^2} + \frac{2}{d} \left( \langle \beta v_2 \rangle - \frac{8\pi^2}{d^2} \|v_2\|_{L_2(\mathbb{R})}^2 \right) + \mathcal{O}(\mu_2 - \alpha_0), \\ \frac{16\pi^2 \langle \beta v_2 \rangle}{(\alpha_0^2 - \mu_2^2)d^3} &= \frac{4\pi \langle \beta v_2 \rangle}{(\alpha_0 - \mu_2)d^2} - \frac{\langle \beta v_2 \rangle}{d} + \mathcal{O}(\mu_2 - \alpha_0). \end{aligned}$$

Thus, to satisfy (6.15), it is sufficient to check that

$$3\langle \beta v_2 \rangle - \frac{16\pi^2}{d^2} \|v_2\|_{L_2(\mathbb{R})}^2 > 0,$$

which is in view of (6.14) equivalent to

$$3\|v_2'\|_{L_2(\mathbb{R})}^2 > \frac{7\pi^2}{d^2} \|v_2\|_{L_2(\mathbb{R})}^2.$$

It is clear that there exists a function  $v \in C_0^\infty(\mathbb{R})$  for which this inequality is valid. Letting  $v_2 := v$  and  $\beta := -v'' + 3\mu_0^2 v$ , we conclude that there exists  $\beta$  such that the inequality (2.19) holds true, if  $\alpha_0$  is sufficiently close to  $\mu_2$  and greater than this number.

## Acknowledgment

The authors are grateful to Miloslav Znojil for many valuable discussions. They also thank the referee for helpful remarks and suggestions.

The work was partially supported by the Czech Academy of Sciences and its Grant Agency within the projects IRP AV0Z10480505 and A100480501, and by the project LC06002 of the Ministry of Education, Youth and Sports of the Czech Republic. D.B. was also supported by RFBR (07-01-00037), by Marie Curie International Fellowship within 6th European Community Framework Programme (MIF1-CT-2005-006254); in addition he gratefully acknowledges the support from Deligne 2004 Balzan Prize in mathematics and the grant of the President of Russian Federation for young scientists and their supervisors (MK-964.2008.1) and for Leading Scientific Schools (NSh-2215.2008.1). D.K. was also supported by FCT, Portugal, through the grant SFRH/BPD/11457/-2002.

## References

- [1] R. A. Adams, *Sobolev spaces*, Academic Press, New York, 1975.
- [2] C. M. Bender, *Making sense of non-Hermitian Hamiltonians*, Rep. Prog. Phys. **70** (2007), 947–1018.
- [3] C. M. Bender and P. N. Boettcher, *Real spectra in non-Hermitian Hamiltonians having  $\mathcal{PT}$  symmetry*, Phys. Rev. Lett. **80** (1998), no. 24, 5243–5246.
- [4] D. Borisov, *Discrete spectrum of a pair of non-symmetric waveguides coupled by a window*, Sbornik Mathematics **197** (2006), no. 4, 475–504.
- [5] D. Borisov, P. Exner, R. Gadyl'shin, and D. Krejčířík, *Bound states in weakly deformed strips and layers*, Ann. H. Poincaré **2** (2002), 553–572.
- [6] W. Bulla, F. Gesztesy, W. Renger, and B. Simon, *Weakly coupled bound states in quantum waveguides*, Proc. Amer. Math. Soc. **125** (1997), 1487–1495.
- [7] E. Caliceti, F. Cannata, and Graffi, *Perturbation theory of  $\mathcal{PT}$ -symmetric Hamiltonians*, J. Phys. A **39** (2006), 10019–10027.
- [8] E. Caliceti, S. Graffi, and J. Sjöstrand, *Spectra of  $\mathcal{PT}$ -symmetric operators and perturbation theory*, J. Phys. A **38** (2005), 185–193.
- [9] P. Dorey, C. Dunning, and R. Tateo, *Spectral equivalences, Bethe ansatz equations, and reality properties in  $\mathcal{PT}$ -symmetric quantum mechanics*, J. Phys. A **34** (2001), 5679–5704.
- [10] P. Duclos and P. Exner, *Curvature-induced bound states in quantum waveguides in two and three dimensions*, Rev. Math. Phys. **7** (1995), 73–102.
- [11] D. E. Edmunds and W. D. Evans, *Spectral theory and differential operators*, Oxford University Press, New York, 1987.
- [12] P. Exner and P. Šeba, *Bound states in curved quantum waveguides*, J. Math. Phys. **30** (1989), 2574–2580.
- [13] R. Gadyl'shin, *On regular and singular perturbation of acoustic and quantum waveguides*, Comptes Rendus Mechanique **332** (2004), no. 8, 647–652.
- [14] I. M. Glazman, *Direct methods of qualitative spectral analysis of singular differential operators*, Israel Program for Scientific Translations, Jerusalem, 1965.
- [15] H.-Ch. Kaiser, H. Neidhardt, and J. Rehberg, *Density and current of a dissipative Schrödinger operator*, J. Math. Phys. **43** (2002), 5325–5350.

- [16] ———, *Macroscopic current induced boundary conditions for Schrödinger-type operators*, Integral Equations and Operator Theory **45** (2003), 39–63.
- [17] ———, *On one-dimensional dissipative Schrödinger-type operators, their dilation and eigenfunction expansions*, Math. Nachr. **252** (2003), 51–69.
- [18] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, Berlin, 1966.
- [19] D. Krejčířík, *Calculation of the metric in the Hilbert space of a  $\mathcal{PT}$ -symmetric model via the spectral theorem*, J. Phys. A: Math. Theor. **41** (2008) 244012.
- [20] D. Krejčířík, H. Břila, and M. Znojil, *Closed formula for the metric in the Hilbert space of a  $\mathcal{PT}$ -symmetric model*, J. Phys. A **39** (2006), 10143–10153.
- [21] D. Krejčířík and J. Kríž, *On the spectrum of curved quantum waveguides*, Publ. RIMS, Kyoto University **41** (2005), no. 3, 757–791.
- [22] H. Langer and Ch. Tretter, *A Krein space approach to  $\mathcal{PT}$ -symmetry*, Czech. J. Phys. **54** (2004), 1113–1120.
- [23] V. P. Mikhailov, *Partial differential equations*, Nauka, Moscow, 1976, in Russian.
- [24] A. Mostafazadeh, *Krein-space formulation of  $\mathcal{PT}$  symmetry,  $\mathcal{CPT}$ -inner products, and pseudo-Hermiticity*, Czech. J. Phys. **56** (2006), 919–933.
- [25] M. Reed and B. Simon, *Methods of modern mathematical physics, IV. Analysis of operators*, Academic Press, New York, 1978.
- [26] K. C. Shin, *On the reality of the eigenvalues for a class of  $\mathcal{PT}$ -symmetric oscillators*, Commun. Math. Phys. **229** (2002), 543–564.
- [27] F. Wolf, *On the essential spectrum of partial differential boundary problems*, Comm. Pure Appl. Math. **12** (1959), 211–228.
- [28] M. Znojil,  *$\mathcal{PT}$ -symmetric square well*, Phys. Lett. A **285** (2001), 7–10.